

Gluconeogenesis and the pentose cycle impact deuterium depleted water efficacy in anticancer therapeutics


László G. Boros¹⁻³, Howard E. Katz⁴, Justine P. Roth⁵, Gábor Somlyai⁶

¹Department of Pediatrics, Harbor-UCLA Medical Center, UCLA School of Medicine, Los Angeles, California, USA, ² The Los Angeles Biomedical Research Institute (LABiOMED), Torrance, California, USA, ³ SIDMAP, LLC, Culver city, California, USA, ⁴Department of Materials Science & Engineering, Whiting School of Engineering, Johns Hopkins University, Baltimore, Maryland, USA., ⁵Departement of Chemistry, Johns Hopkins University, Baltimore, Maryland, USA., ⁶HYD, LLC for Cancer Research & Drug Development, Budapest, Hungary, European Union (EU)

Increased glucose decarboxylation during nucleic acid ribose synthesis occurs at the expense of glycolysis. This pathway is coupled to the recycling of drinking water, which passes through the cytoplasm after absorption. Consumption of water from the cytoplasm and cellular matrix, upon ring-opening hydrolysis in the pentose cycle, results in the production of 6-phospho-D-gluconate. Oxidation at C1 of this substrate, together with NADP⁺ reduction, transforms mammalian cells into megaloblastic and sideroblastic with “unstable” DNA [1-3]. On the other hand, gluconeogenesis is coupled to mitochondrial substrate oxidation and the recycling of “matrix water”, which passes by substrate shuttling during respiration and hydration of metabolites in mitochondria. Results will be presented which suggest that mitochondrial matrix water (re)cycling [4, 5] during oxidation of ketogenic substrates, low in deuterium content [6], reverses the pentose cycle-transformed tumorous phenotype [4]. The anti-cancer drug Avastin® requires mitochondrial complex-I activation with normal mitochondrial morphology [7] for efficacy, which demonstrates the substitution of natural abundance water by deuterium depleted water (DDW) produced in the mitochondrial matrix. Stable isotope depletion is also involved in the efficacy of Glivec®, which is associated with increased fatty acid oxidation through the tricarboxylic acid (TCA) or Krebs cycle [8-10]. This cycle produces energy in the form of adenosine triphosphate (ATP) via the aerobic oxidation of acetate derived from carbohydrates, fats and proteins. In addition to CO₂, the reduced nicotinamide cofactor and precursors of certain amino acids are products. Clearly, the central importance of the TCA cycle in many biochemical pathways makes it a critical target for controlling cellular metabolism, growth and survival. Using DDW in place of natural abundance water and decreasing deuterium by even 1 parts per million (ppm) for every 8 hours in cultures impedes the growth of MIA-PaCa and MCF-7 cancer cell lines (xCELLigence System - Roche). The effect is thought to derive from interfering with the pentose cycle and subsequently altering the structural integrity of cellular DNA in tumor cells. Further, it is proposed that “metabolic” matrix water production, upon ketogenic substrate oxidation, precipitates the DNA strand-breaking event at the accessible surface areas of deoxyribose [11] via gluconeogenesis. It follows from observations, with the anti-cancer drug Rosiglitazone®, that isotope depletion in the TCA cycle and “metabolic” water affects the deuterium content at the C5'-C3' hydrogenic positions (Fig 1-2) of the nucleotide sugar fragment, after fatty acid chain shortening and redistribution of low-deuterium acetyl-CoA [12, 13]. The observations link DDW, containing 100, 50 and 25 ppm deuterium, which is 155 ppm in natural abundance water (non-DDW)) to limited tumor cell growth and extension of median survival time in patients with inoperable advanced pancreatic cancer from 6 months with chemotherapy alone to 40 months, when DDW is added as an adjuvant within 60 days after diagnosis in an open label trial. The clinical and experimental data implicate exchange of pentose cycle-derived triose fragments (C1-C3 to C3-C5) into ribose and deoxyribose [14], and the nucleic acid sugar backbone as a sink for deuterium originating from cytoplasmic water.

These reactions involving carbon and hydrogen bonds give rise to apparent (and potentially complex) isotope effects [15-17], which might be exploited in anti-cancer chemotherapy [18, 19]. Additional applications include intramolecular deuterium disequilibrium as the result of deuterium depleting mechanisms linked with isomerase function [20] as novel NMR based biomarkers to determine risks for aneuploidy, as well as response to various deuterium depleting therapies in disease. Water distribution in tumors in comparison with that of surrounding tissues, by means of magnetic resonance imaging (MRI), is routinely observed by the deuterium oxide (D₂O)/water (H₂O) contrast, where D₂O occupying tumors cause a reduction of the signal when ¹H detection is used during flow velocity studies. Perfusion with D₂O may reduce water signals to minimum, which can subsequently be renewed with distilled low deuterium containing H₂O in various models [23]. Therefore, the replacement of low deuterium-containing metabolic matrix water due to impaired mitochondria with that of more D₂O enriched water from the circulation in the cytoplasm of cancer cells can potentially induce similar losses of the proton signal, whereby the resulting D₂O/H₂O contrast obtained with MRI contributes an effective clinically applied metabolic biomarker tool to studying tumors within various organs.

Figure 1. Comparison of metabolic profile changes associated with 1) natural deuterium depletion by low deuterium fatty acid oxidation. Avastin®, Glivec® and Rosiglitazone® exert similar effect and require intact mitochondria for efficacy [7, 9, 13] (**Red boxes No1**), and 2) low deuterium metabolic water recycling from the mitochondrial matrix during citrate, isocitrate and malate formation; the target of fumarate hydratase activation [4, 5] and hyperbaric oxygen treatment combined with a ketogenic diet [21] (**Red box No2**). Mitochondrial shuttles, such as the malate shuttle, pass low deuterium carrying fatty acid carbons to gluconeogenesis, where glyceraldehyde-3-phosphate becomes the source of extensive carbon exchange reactions [14] for the non-oxidative pentose cycle to maintain low deuterium saturation in C3'-C5' pentose sugar carbon positions (**Red box No3**). These are the carbon sites where DNA stability, radiation- and chemotherapy derived hydroxyl radical sensitivities are regulated by hydrogen/deuterium [11] due to primary and secondary intrinsic isotope effects; as well as partially by proton tunneling [15-17]. Besides the C3'-C5' nucleic acid sugar backbone fragment, *de novo* nucleic acid base syntheses, hydrogen bonding and deuterium channeling into hydrogen bonds are controlled by the serine oxidation glycine cleavage single carbon cycle pathways [22; SOGC] (**Red box No4**). When tumor cells revert to the Warburg phenotype and reductive carboxylation-driven mitochondria, deuterium depletion in free (drinking) water becomes an effective deuterium depleting mechanism for specific carbon sites in nucleic acid backbone sugars and the bases (**Red box No5**). (Blue arabic numbers are enzyme identifiers also found in ref# [12])



References:

1. Green, R. Mystery of thiamine-responsive megaloblastic anemia unlocked. *Blood*. 102, 3464-3465 (2003). doi: 10.1182/blood-2003-09-3025
2. Fleming, J. C., *et al.* Characterization of a murine high-affinity thiamine transporter, Slc19a2. *Mol Genet Metab*. 74, 273-280 (2001). doi:10.1006/mgme.2001.3241
3. Boros, L. G., *et al.* Defective RNA ribose synthesis in fibroblasts from patients with thiamine-responsive megaloblastic anemia (TRMA). *Blood*. 102, 3556-61 (2003). doi: 10.1182/blood-2003-05-1537
4. Tong, W. H. *et al.* The glycolytic shift in fumarate-hydratase-deficient kidney cancer lowers AMPK levels, increases anabolic propensities and lowers cellular iron levels. *Cancer. Cell*. 20, 315-327 (2011).. doi: 10.1016/j.ccr.2011.07.018
5. Yang, Y., *et al.* Metabolic reprogramming for producing energy and reducing power in fumarate hydratase null cells from hereditary leiomyomatosis renal cell carcinoma. *PLoS. One*. 8, e72179 (2013). doi: 10.1371/journal.pone.0072179
6. Duan, J. R., *et al.* Natural deuterium distribution in fatty acids isolated from peanut seed oil: a site-specific study by quantitative ²H NMR spectroscopy. *Chembiochem*. 3, 752-759 (2002). doi: 10.1002/1439-7633(20020802)
7. Xu, J., *et al.* Colorectal cancer cells refractory to anti-VEGF treatment are vulnerable to glycolytic blockade due to persistent impairment of mitochondria. *Mol Cancer Ther*. 12, 717-24 (2013). doi: 10.1158/1535-7163.MCT-12-1016-T
8. Boros, L. G., *et al.* Imatinib and chronic-phase leukemias. *N. Engl. J. Med*. 347, 67-68 (2002). doi: 10.1056/NEJM200207043470116
9. Kominsky, D. J., *et al.* Abnormalities in glucose uptake and metabolism in imatinib-resistant human BCR-ABL-positive cells. *Clin. Cancer. Res*. 15, 3442-50 (2009). doi: 10.1158/1078-0432.CCR-08-3291
10. Boren, J. *et al.* Gleevec (STI571) influences metabolic enzyme activities and glucose carbon flow toward nucleic acid and fatty acid synthesis in myeloid tumor cells. *J. Biol. Chem*. 276, 37747-37753 (2001). doi: 10.1074/jbc.M105796200
11. Balasubramanian, B., *et al.* DNA strand breaking by the hydroxyl radical is governed by the accessible surface areas of the hydrogen atoms of the DNA backbone. *Proc. Natl. Acad. Sci. USA*. 95, 9738-9743 (1998). doi: 10.1073/pnas.95.17.9738
12. Boros L. G., *et al.* Targeted ¹³C-labeled tracer fate associations for drug efficacy testing in cancer. *In: Tumor Cell Metabolism – Pathways, Regulation & Biology*, Eds. Shoshan, M. (Karolinska Institute, Stockholm, Sweden) & Mazurek, S. L. (Justus-Liebig-University, Giessen, Germany), Chapter 15 - pages 349-370 (2014). ISBN 978-3-7091-1823-8 ISBN 978-3-7091-1824-5 (eBook), Springer Wien Heidelberg New York Dordrecht London. doi 10.1007/978-3-7091-1824-5
13. Harrigan, G. G. *et al.* PNU-91325 increases fatty acid synthesis from glucose and mitochondrial long chain fatty acid degradation: a comparative tracer-based metabolomics study with rosiglitazone and pioglitazone in HepG2 cells. *Metabolomics*. 2, 21-29 (2006). doi: 10.1007/s11306-006-0015-5

14. Centelles, J. J., *et al.* Metabolic profile and quantification of deoxyribose synthesis pathways in HepG2 cells. *Metabolomics*. 3, 105-111 (2007). doi: 10.1007/s11306-006-0049-8
15. Löwdin, P-O. Proton Tunneling in DNA and its Biological Implications. *Rev. Mod. Phys.* 35, 724 (1963). doi: 10.1103/RevModPhys.35.724
16. Sobczyk, L., *et al.* H/D Isotope Effects in Hydrogen Bonded Systems. *Molecules*. 18, 4467-4476 (2013). doi:10.3390/molecules18044467
17. Drechsel-Grau, C., *et al.* Exceptional isotopic-substitution effect: breakdown of collective proton tunneling in hexagonal ice due to partial deuteration. *Angew Chem Int Ed Engl.* 53, 10937-10940 (2014). doi: 10.1002/anie.201405989
18. Boros, L. G., *et al.* Fumarate hydratase and deuterium depletion control oncogenesis via NADPH-dependent reductive synthesis: mitochondrial matrix water, DNA deuteration and epigenetic events. *In: Proceedings of the 105th Annual Meeting of the American Association for Cancer Research; 2014 Apr 5-9; San Diego, CA. Philadelphia (PA): AACR; Cancer Res 2014; 74(19 Suppl):Abstract nr 1426.* doi:10.1158/1538-7445.AM2014-1426
19. Somlyai, G., *et al.* Naturally occurring deuterium is essential for the normal growth rate of cells. *FEBS Lett.* 317, 1–4 (1993). doi: 10.1016/0014-5793(93)81479-J
20. Schleucher, J., *et al.* Intramolecular deuterium distributions reveal disequilibrium of chloroplast phosphoglucose isomerase. *Plant, Cell & Environment* 22, 525–533 (1999). doi: 10.1046/j.1365-3040.1999.00440
21. Poff, A., *et al.* The ketogenic diet and hyperbaric oxygen therapy prolong survival in mice with systemic metastatic cancer. *PLoS. One.* 8, e65522 (2013). doi: 10.1371/journal.pone.0065522
22. Tedeschi, P. M., *et al.* Contribution of serine, folate and glycine metabolism to the ATP, NADPH and purine requirements of cancer cells. *Cell. Death. Dis.* 4, e877 (2013). doi: 10.1038/cddis.2013.393
23. Ilvonena, K., *et al.* MRI-Based D₂O/H₂O-Contrast Method to Study Water Flow and Distribution in Heterogeneous Systems: Demonstration in Wood Xylem. *J. Magn. Reson.* 149, 36–44 (2001). doi: 10.1006/jmre.2001.2284

Author Information:

Correspondence and requests for materials should be addressed to L.G.B. (boros@labiomed.org; boros.laszlo@yahoo.com) (USA)

Key Words:

deuterium depletion, mitochondrial matrix metabolic water, deuterium-depleted water, NADPH, oxidative pentose cycle, median survival time, unresectable pancreatic cancer, open label study, targeted tracer fate associations study, targeted deuterium fate association study, TTFAS, TDFAS

Additional ¹³C tracer and data interpretations can be found on the World Wide Web at <http://youtu.be/Pms6k9AQ3bQ> regarding mammalian cell mitochondria.

DOI: 10.13140/2.1.3536.9281